arXiv:1605.06583v1 [math.RA] 21 May 2016

BREADTH AND CHARACTERISTIC SEQUENCE OF NILPOTENT LIE ALGEBRAS

ELISABETH REMM

Abstract. The notion of breadth of a nilpotent Lie algebra was introduced and used to approach problems of classification up to isomorphism in [6]. In the present paper, we study this invariant in terms of characteristic sequence, another invariant introduced by Goze and Ancochea in [2]. This permits to complete the determination of Lie algebras of breadth 2 studied in [6] and to begin the work for Lie algebras with breadth greater than 2.

Introduction

The breadth of a Lie algebra was introduced in [6]. As it is well explained in this paper, initially, the notion of breadth was defined to investigate the p-groups: the breadth of a p-group is the size of the largest conjugacy class in this p-group. Recently characterizations of finite p-groups of breadth 1 and 2 have been done. In [6], the authors translate this notion to finite dimensional nilpotent Lie algebras: the breadth of a nilpotent Lie algebra is the maximum of the ranks of the adjoint operators and in their paper, they describe nilpotent Lie algebras of breadth 1 and, for dimensions 5 and 6, of breadth 2.

We know that the classification, up to isomorphism, of finite dimensional nilpotent Lie algebras is a very difficult problem. It is solved up to the dimension 7 in the complex and real cases and for some special families of nilpotent Lie algebras in bigger dimensions. In recent papers [8, 9], since the classification problem in large dimensions seems unrealistic, we choose to describe classes of nilpotent Lie algebras, in any dimension, associated with the characteristic sequence. For these families, we have reduced the notion of Chevalley-Eilenberg cohomology and given, sometimes, topological properties of these families. In this paper, we consider these results to complete in any dimension the characterization of nilpotent Lie algebras of breadth equal to 2, completing, in some way, the work [6]. We develop also the case where the breadth is 3.

1. breadth and characteristic sequence

1.1. Breadth of a nilpotent Lie algebra.

Let $\mathfrak{g}$ be a finite dimensional Lie algebra over field $\mathbb{K}$ of characteristic 0.

Definition 1. [6] *The breadth of $\mathfrak{g}$ is the invariant, up to isomorphism,*

$$b(\mathfrak{g}) = \max\{\text{rank}(\text{ad}X),\ X \in \mathfrak{g}\}.$$

More generally, if I is a not trivial ideal of $\mathfrak{g}$, then the breadth of $\mathfrak{g}$ on I is

$$b_I(\mathfrak{g}) = \max\{\text{rank}(\text{ad}X|_I),\ X \in \mathfrak{g}\}.$$

Date: Chevat 16, 5775.

We have $b_{\mathfrak{g}}(\mathfrak{g}) = b(\mathfrak{g})$ and clearly $b_I(\mathfrak{g}) \leq b(\mathfrak{g})$. For example, $\mathfrak{g}$ is abelian if and only if $b(\mathfrak{g}) = 0$ and $\mathfrak{g}$ is filiform (see [5]) if and only if $b(\mathfrak{g}) = \dim \mathfrak{g} - 2$. Recall that a finite dimensional nilpotent Lie algebra is called filiform, notion introduced by Michele Vergne, if $\mathfrak{g}$ satisfies

$$\dim \mathcal{C}_i(\mathfrak{g}) = i$$

for $i = 0, \cdots, n-2$ where $n = \dim \mathfrak{g}$ and $\mathcal{C}_i(\mathfrak{g})$ are the terms of the ascending sequence of $\mathfrak{g}$ that is

$$\begin{cases} \mathcal{C}_0(\mathfrak{g}) = \{0\}, \\ \mathcal{C}_i(\mathfrak{g}) = \{X \in \mathfrak{g}, [X, \mathfrak{g}] \in \mathcal{C}_{i-1}(\mathfrak{g})\}. \end{cases}$$

The following proposition is proved in [6]

Proposition 2. *For any finite dimensional Lie algebra $\mathfrak{g}$, we have*

$$b(\mathfrak{g}) \leq \dim(\mathfrak{g}/Z(\mathfrak{g})) - 1.$$

In the case of a filiform Lie algebra, the upper bound of this inequality is satisfied. In fact if $\mathfrak{g}$ is filiform, then $\dim Z(\mathfrak{g}) = 1$ and $b(\mathfrak{g}) = n - 2 = \dim(\mathfrak{g}/Z(\mathfrak{g})) - 1$.

Assume now that $\mathfrak{g}$ is finite dimensional and nilpotent. We have seen that $b(\mathfrak{g}) = 0$ if and only if $\mathfrak{g}$ is abelian. Now if $b(\mathfrak{g}) = 1$, then, from [6] we have $\dim \mathcal{C}^1(\mathfrak{g}) = 1$ where $\mathcal{C}^1(\mathfrak{g})$ is the derived subalgebra of $\mathfrak{g}$, that is generated by the Lie brackets $[X, Y]$ for any $X, Y \in \mathfrak{g}$. Then, if $\mathfrak{g}$ is indecomposable, that is without direct factor, then $\mathfrak{g}$ is an Heisenberg Lie algebra. The non trivial study begins with $b(\mathfrak{g}) = 2$. This is a large part of the work of [6]. We will complete their result using and comparing with an another invariant of nilpotent Lie algebras, the characteristic sequence, presented in the following section.

1.2. **Characteristic sequence.** Let $\mathfrak{g}$ be a finite dimensional nilpotent Lie algebra over $\mathbb{K}$. Let $\mathcal{C}^1(\mathfrak{g})$ and $Z(\mathfrak{g})$ be respectively the derived subalgebra and center of $\mathfrak{g}$. The characteristic sequence of $\mathfrak{g}$, sometimes called the Goze's invariant and introduced in [2], is an useful invariant in the classification theory of finite-dimensional nilpotent Lie algebras. It is define as follows: for any $X \in \mathfrak{g}$, let $c(X)$ be the ordered sequence, for the lexicographic order, of the dimensions of the Jordan blocks of the nilpotent operator adX. The characteristic sequence of $\mathfrak{g}$ is the invariant, up to isomorphism,

$$c(\mathfrak{g}) = \max\{c(X),\ X \in \mathfrak{g} - \mathcal{C}^1(g)\}.$$

Then $c(\mathfrak{g})$ is a sequence of type $(c_1, c_2, \cdots, c_k, 1)$ with $c_1 \geq c_2 \geq \cdots \geq c_k \geq 1$ and $c_1 + c_2 + \cdots + c_k + 1 = n = \dim \mathfrak{g}$. For example,

(1) $c(\mathfrak{g}) = (1, 1, \cdots, 1)$ if and only if $\mathfrak{g}$ is abelian,
(2) $c(\mathfrak{g}) = (2, 1, \cdots, 1)$ if and only if $\mathfrak{g}$ is a direct product of an Heisenberg Lie algebra by an abelian ideal,
(3) If $\mathfrak{g}$ is 2-step nilpotent then there exists p and q such that $c(\mathfrak{g}) = (2, 2, \cdots, 2, 1, \cdots, 1)$ with $2p + q = n$, that is p is the occurrence of 2 in the characteristic sequence and q the occurrence of 1.
(4) $\mathfrak{g}$ is filiform if and only if $c(\mathfrak{g}) = (n-1, 1)$.

A vector $X \in \mathfrak{g} - \mathcal{C}^1(\mathfrak{g})$ is called a characteristic vector if $c(X) = c(\mathfrak{g})$. Of course, such a vector is not unique.

1.3. **Breadth and characteristic sequence.** Let $\mathfrak{g}$ be a finite dimensional nilpotent Lie algebra, $b(\mathfrak{g})$ its breadth and $c(\mathfrak{g})$ its characteristic sequence.

Lemma 3. *Let X be a characteristic vector with $c(\mathfrak{g}) = c(X) = (c_1, \cdots, c_k, 1)$. Then*

$$\operatorname{rank}(\operatorname{ad}X) = \sum_{i=1}^{k} c_i - k \tag{1}$$

In fact, if we consider the Jordan block J_i of dimension c_i of the operator $\operatorname{ad}X$, since 0 is the only eigenvalue, we have $\operatorname{rank}(J_i) = c_i - 1$.

We deduce

Theorem 4. *Let $\mathfrak{g}$ be a finite dimensional Lie algebra whose characteristic sequence is $c(\mathfrak{g}) = (c_1, \cdots, c_k, 1)$. Then, its breadth satisfies*

$$b(\mathfrak{g}) = \sum_{i=1}^{k} c_i - k. \tag{2}$$

Consequences.

1. If $\mathfrak{g}$ is n-dimensional 2-step nilpotent, then $c(\mathfrak{g}) = (2, \cdots, 2, 1, \cdots, 1)$. If 2 appears p times, then

$$b(\mathfrak{g}) = p.$$

2. If $\mathfrak{g}$ is n-dimensional 3-step nilpotent, then $c(\mathfrak{g}) = (3, \cdots, 3, 2, \cdots, 2, 1, \cdots, 1)$. If 3 appears p_3 times, 2 appears p_2 times in this sequence, then

$$b(\mathfrak{g}) = 2p_3 + p_2.$$

3. More generally, if $\mathfrak{g}$ is n-dimensional k-step nilpotent, then its characteristic sequence is $c(\mathfrak{g}) = (k, \cdots, k, \cdots, 3, \cdots, 3, 2, \cdots, 2, 1, \cdots, 1)$. Assume that k appears p_k-times, $k-1$ p_{k-1}-times, up to 2 which appears p_2-times and 1 which appears p_1-times. Then

$$b(\mathfrak{g}) = (k-1)p_k + (k-2)p_{k-1} + \cdots + 2p_3 + p_2 = \sum_{i=1}^{k} (i-1)p_i.$$

Proposition 5. *A n-dimensional nilpotent Lie algebra satisfies $b(\mathfrak{g}) = 2$ if and only if one of these conditions is satisfied:*

1. *$\mathfrak{g}$ is 2-step nilpotent with $c(\mathfrak{g}) = (2, 2, 1, \cdots, 1)$.*
2. *$\mathfrak{g}$ is 3-step nilpotent with $c(\mathfrak{g}) = (3, 1, \cdots, 1)$.*

In fact, if $\mathfrak{g}$ is k-step nilpotent with $k \geq 4$, then $b(\mathfrak{g}) = (k-1)p_k + (k-2)p_{k-1} + \cdots + 2p_3 + p_2$ with $p_k > 0$. Then $b(\mathfrak{g}) \geq 3$. Now if $\mathfrak{g}$ is 3-step nilpotent, then $b(\mathfrak{g}) = 2p_3 + p_2$ with $p_3 > 0$. Then $b(\mathfrak{g}) = 2$ implies $p_3 = 1$ and $p_2 = 0$. If $\mathfrak{g}$ is 2-step nilpotent, then $b(\mathfrak{g}) = p_2$ this gives $p_2 = 2$.

2. Nilpotent Lie algebras with breadth equal to 2

A lot of results concerning nilpotent Lie algebras with breadth equal to 2 are presented in [6], principally for the small dimensions. In this section, we propose to complete this study using Proposition 5. But, before begining this study, we recall some properties of an important tool used in [8] and based on deformation theory of Lie algebras. Let $\mathfrak{g}_0 = (\mathbb{K}^n, \mu_0)$ be a n-dimensional Lie algebra over $\mathbb{K}$ where μ_0 denotes the Lie bracket of $\mathfrak{g}_0$ and $\mathbb{K}^n$ the underlying linear space to $\mathfrak{g}_0$. A formal deformation $\mathfrak{g}$ of $\mathfrak{g}_0$ is given by a pair $\mathfrak{g} = (\mathbb{K}^n, \mu)$ where μ is a formal series

$$\mu = \mu_0 + \sum t^i \varphi_i$$

with skew-symmetric bilinear maps φ_i on $\mathbb{K}^n$ and μ satisfying the Jacobi identity. To simplify notations, we denote for any bilinear map φ on $\mathbb{K}^n$ with values on $\mathbb{K}^n$, $\varphi \bullet \varphi$ the trilinear map

$$\varphi \bullet \varphi(X,Y,Z) = \varphi(\varphi(X,Y),Z) + \varphi(\varphi(Y,Z),X) + \varphi(\varphi(Z,X),Y)$$

and by $\varphi \circ_1 \varphi$ the comp_1-operation (the composition on the first element), that is

$$\varphi \circ_1 \varphi(X,Y,Z) = \varphi(\varphi(X,Y),Z).$$

Then the Jacobi identity for φ is equivalent to $\varphi \bullet \varphi = 0$. If μ is a formal deformation of μ_0, then $\mu \bullet \mu = 0$ implies

$$\begin{cases} \delta_{C,\mu_0}\varphi_1 = 0, \\ \varphi_1 \bullet \varphi_1 = \delta_{C,\mu_0}\varphi_2, \end{cases}$$

where δ_{C,μ_0} is the coboundary operator associated to the Chevalley-Eilenberg complex of $\mathfrak{g}_0$:

$$\delta_{C,\mu_0}(\varphi) = \mu_0 \bullet \varphi + \varphi \bullet \mu_0.$$

We shall denote by $H^*_C(\mathfrak{g}_0, \mathfrak{g}_0)$ or sometimes $H^*_C(\mu_0, \mu_0)$ the corresponding cohomology spaces. A special class of formal deformation is the class of linear deformations. We call *linear deformation* of $\mathfrak{g}_0 = (\mathbb{K}^n, \mu_0)$ a formal deformation $\mu = \mu_0 + \sum t^i \varphi_i$ with $\varphi_k = 0$ for any $k \geq 2$, that is

$$\mu = \mu_0 + t\varphi_1.$$

In this case, the Jacobi identity for μ is equivalent to

$$\begin{cases} \delta_{C,\mu_0}\varphi_1 = 0, \\ \varphi_1 \bullet \varphi_1 = 0. \end{cases} \tag{3}$$

2.1. **Lie algebras with characteristic sequence** $(3, 1, \cdots, 1)$**.** Let $\mathfrak{g} = (\mathbb{K}^n, \mu)$ be a n-dimensional nilpotent Lie algebra with characteristic sequence $c(\mathfrak{g}) = (3, 1, \cdots, 1)$. Then from [8] it is isomorphic to a linear deformation of the Lie algebra $\mathfrak{g}_{1,0,n-1} = (\mathbb{K}^n, \mu_0)$ whose Lie brackets are

$$[X_1, X_2] = X_3,\ [X_1, X_3] = X_4,\ [X_1, X_i] = 0 \text{ for } i = 4, \cdots, n,\ [X_i, X_j] = 0,\ 2 \leq i < j \leq n.$$

So $\mu = \mu_0 + t\varphi$ where φ is a bilinear map satisfying the conditions (3) and conditions which imply that μ is a Lie bracket of a 3-step nilpotent Lie algebra. All these conditions are described in [8] and imply that $\mathfrak{g}$ is in one of the following cases:

(1) $\mathfrak{g}$ is isomorphic to $\mathfrak{g}_{1,0,n-1}$,
(2) $\mathfrak{g}$ is a direct product of an abelian ideal and of the 5-dimensional Lie algebra and its Lie bracket is isomorphic to $\mu_0 + \varphi$ with $\varphi(X_2, X_3) = aX_5$,

(3) $\mathfrak{g}$ is a direct product of an abelian ideal and of the 5-dimensional Lie algebra and its Lie bracket is isomorphic to $\mu_0 + \varphi$ with $\varphi(X_2, X_5) = bX_4$
(4) If $\mathfrak{g}$ is indecomposable and of dimension greater than 5, then $\mathfrak{g}$ is isomorphic to the Lie algebra

$$\begin{cases} [X_1, X_2] = X_3,\ [X_1, X_3] = X_4, \\ [X_2, X_k] = a_{2,k}X_4,\ k \geq 5, \\ [X_l, X_k] = a_{l,k}X_4,\ 5 \leq l < k \leq n. \end{cases}$$

This last family of Lie algebras, parametrized by the constants structure $a_{2,k}$ and $a_{l,k}$ can be reduced to obtain a classification up to isomorphism.

Proposition 6. [4] *Any n-dimensional nilpotent Lie algebra, $n \geq 4$ with breadth equal to 2 and characteristic sequence $(3, 1, \cdots, 1)$ is isomorphic to one of the following Lie algebras whose Lie brackets are*

(1) $[X_1, X_2] = X_3,\ [X_1, X_3] = X_4$ that is $\mathfrak{g}_{1,0,n-1}$.
(2) $[X_1, X_2] = X_3,\ [X_1, X_3] = X_4, [X_2, X_3] = X_5$, here $n \geq 5$.
(3) $[X_1, X_2] = X_3,\ [X_1, X_3] = X_4, [X_{2i+1}, X_{2i+2}] = X_4, i = 1, \cdots, p-1$ with $n = 2p$,
(4) $[X_1, X_2] = X_3,\ [X_1, X_3] = X_4, [X_2, X_n] = X_4, [X_{2i+1}, X_{2i+2}] = X_4, i = 1, \cdots, p-1$ with $n = 2p+1$,

2.2. **Lie algebras whose characteristic sequence is $(2, 2, 1, \cdots, 1)$.** In [7], we have studied nilpotent Lie algebras with characteristic sequence $(2, 2, 1, 1, 1, 1)$ in order to construct symplectic form on these Lie algebras. We will use notations and results which are in this paper. Let $\mathcal{F}^{n,2}$ be the set of n-dimensional 2-step nilpotent Lie algebras and $\mathcal{F}^{n,2}_{2,n-2}$ the subset corresponding to Lie algebras of characteristic sequence $(2, 2, 1, \cdots, 1)$ (the subscripts correspond to the numbers of 2 and 1 in the characteristic sequence). Let $\mathfrak{g}_{2,n-2}$ be the n-dimensional Lie algebra given by the brackets

$$[X_1, X_{2i}] = X_{2i+1}, \quad i = 1, 2.$$

It is obvious that $\mathfrak{g}_{2,n-2} \in \mathcal{F}^{n,2}_{2,n-2}$. From [7], any n-dimensional 2-step nilpotent Lie algebra with characteristic sequence $(2, 2, 1, \cdots, 1)$ is isomorphic to a linear deformation of $\mathfrak{g}_{2,n-2}$. This means that its Lie bracket is isomorphic to $\mu = \mu_0 + t\varphi$, where μ_0 is the Lie bracket of $\mathfrak{g}_{2,n-2}$, and φ is a skew-bilinear map such that

$$\begin{cases} \varphi \in Z^2_{CH}(\mathfrak{g}_{2,n-2}, \mathfrak{g}_{2,n-2}), \\ \varphi \circ_1 \varphi = 0. \end{cases}$$

Recall that $Z^2_{CH}(\mathfrak{g}_{2,n-2}, \mathfrak{g}_{2,n-2})$ is constituted of bilinear maps φ on $\mathbb{K}^n$ with values in this space such that

$$\varphi \circ_1 \mu_0 + \mu_0 \circ_1 \varphi = 0.$$

In fact, in [8], we have define a sub-complex of the Chevalley-Eilenberg complex whose main property is that for any deformation $\mu = \mu_0 + \sum t^i \varphi_i$ of a 2-step nilpotent Lie algebra $\mathfrak{g}_0 = (\mu_0, \mathbb{K}^n)$ which is also 2-step nilpotent, then $\varphi_1 \in Z^2_{CH}(\mu_0, \mu_0)$.

We deduce that the family $\mathcal{F}^{n,2}_{2,n-2}$ of n-dimensional 2-step nilpotentLie algebras with characteristic sequence $(2, 2, 1, \cdots, 1)$ is the union of two algebraic components, the first one,

$\mathcal{C}_1(\mathcal{F}^{n,2}_{2,n-2})$, corresponding to the cocycles

$$\varphi(X_i, X_j) = \sum_{k=1}^{2} a_{i,j}^{2k+1} X_{2k+1}, \quad 2 \leq i < j \leq n, \quad i,j \neq 3,5 \tag{4}$$

the second one, $\mathcal{C}_2(\mathcal{F}^{n,2}_{2,n-2})$, to the cocyles

$$\begin{cases} \varphi(X_i, X_j) = \sum_{k=1}^{2} a_{i,j}^{2k+1} X_{2k+1}, \quad (i,j) \neq (2,4),\ 2 \leq i < j \leq n-1, \quad i,j \notin \{3,5\} \\ \varphi(X_2, X_4) = X_n \end{cases} \tag{5}$$

where the undefined products $\varphi(X,Y)$ are nul or obtained by skew-symmetry. Each of these components is a regular algebraic variety. These components can be characterized by the number of generators.

Proposition 7. *Any n-dimensional nilpotent Lie algebra with breadth equal to 2 and characteristic sequence $(2,2,1,\cdots,1)$ is isomorphic to a Lie algebra belonging to one of the components $\mathcal{C}_1(\mathcal{F}^{n,2}_{2,n-2})$ or $\mathcal{C}_2(\mathcal{F}^{n,2}_{2,n-2})$.*

3. Nilpotent Lie algebras with breadth equal to 3

From Theorem 4, the characteristic sequence of a n-dimensional Lie algebra of breadth $b(\mathfrak{g}) = 3$ is equal to one of the following:

(1) $c(\mathfrak{g}) = (4,1,\cdots,1)$, $(n \geq 5)$,
(2) $c(\mathfrak{g}) = (3,2,1,\cdots,1)$, $(n \geq 6)$,
(3) $c(\mathfrak{g}) = (2,2,2,1,\cdots,1)$, $(n \geq 7)$.

3.1. **Nilpotent Lie algebras of characteristic sequence** $(4,1,\cdots,1)$**.** Let us consider the n-dimensional nilpotent Lie algebra $\mathfrak{g}_{1,0,0,n-4} = (\mathbb{K}^n, \mu_0)$ with characteristic sequence $(4,1,\cdots,1)$ whose Lie brackets in the basis $\{X_1,\cdots,X_n\}$ are

$$\mu_0(X_1, X_i) = X_{i+1},\ i = 2,3,4$$

all the others brackets being nul. Any n-dimensional Lie algebra $\mathfrak{g}$ whose characteristic sequence is also $(4,1,\cdots,1)$ is isomorphic to a Lie algebra whose Lie bracket μ is a linear deformation of μ_0, that is $\mu = \mu_0 + t\varphi$ where φ is a 2-cocycle of the Chevalley-Eilenberg complex $H^*_C(\mu_0,\mu_0)$ satisfying

$$\begin{cases} \delta^2_{C,\mu_0}(\varphi) = 0, \\ \varphi \bullet \varphi = 0, \\ \delta^2_{R,\mu_0}(\varphi) = 0, \\ \mu_0^2 \circ_1 \varphi^2 + \mu_0 \circ_1 \varphi \circ_1 \mu_0 \circ_1 \varphi + \mu_0 \circ_1 \varphi^2 \circ_1 \mu_0 + \varphi \circ_1 \mu_0^2 \circ_1 \varphi + \\ + \varphi \circ_1 \mu_0 \circ_1 \varphi \circ_1 \mu_0 + \varphi 2 \circ_1 \mu_0^2 = 0, \\ \mu_0 \circ_1 \varphi^3 + \varphi \circ_1 \mu_0 \circ_1 \varphi^2 + \varphi^2 \circ_1 \mu_0 \circ_1 \varphi + \varphi^3 \circ_1 \mu_0 = 0, \\ \varphi^4 = 0, \end{cases} \tag{6}$$

where $\delta^2_{R,\mu_0}(\varphi)$ is the 5-linear map

$$\delta^2_{R,\mu_0}(\varphi) = \mu_0^3 \circ_1 \varphi + \mu_0^2 \circ_1 \varphi \circ_1 \mu_0 + \mu_0 \circ_1 \varphi \circ_1 \mu_0^2 + \varphi \circ_1 \mu_0^3,$$

and $\varphi^k = \varphi \circ_1 \cdots \circ_1 \varphi$. Moreover, φ satisfies $\varphi(X_1, Y) = 0$ for any Y and X_1 is not in the derived subalgebra of $\mathfrak{g}$. This implies that $(\mu_0 \circ_1 \varphi)(X_i, X_j, X_k) = 0$ if $1 < i < j < k$. Then (6) gives

$$(7) \qquad \begin{cases} \varphi(X_5, X_i) = 0, \ \forall i, \\ \varphi(X_2, X_3) = aX_5 + \sum_{k=6}^{n} d_k X_k, \\ \varphi(X_2, X_k) = a_k X_4 + b_k X_5 \ \ 6 \leq k \leq n, \\ \varphi(X_3, X_k) = a_k X_5 \ \ 6 \leq k \leq n, \\ \varphi(X_i, X_j) = c_{i,j} X_5 \ \ 6 \leq i < j \leq n. \end{cases}$$

Proposition 8. *Every n-dimensional nilpotent Lie algebra ($n \geq 5$) of breadth 3 and characteristic sequence $(4, 1, \cdots, 1)$ is isomorphic to a Lie algebra belonging to one of the families*

$$\begin{cases} [X_1, X_i] = X_{i+1}, \quad i = 2, 3, 4, \\ [X_5, X_i] = 0, \ \forall i, \\ [X_2, X_3] = aX_5, \\ [X_2, X_k] = a_k X_4 + b_k X_5, \quad 6 \leq k \leq n, \\ [X_3, X_k] = a_k X_5, \quad 6 \leq k \leq n, \\ [X_i, X_j] = c_{i,j} X_5, \quad 6 \leq i < j \leq n. \end{cases}$$

$$\begin{cases} [X_1, X_i] = X_{i+1}, \quad i = 2, 3, 4, \\ [X_5, X_i] = 0, \ \forall i \\ [X_2, X_3] = aX_5 + \sum_{k=6}^{n} d_k X_k, \\ [X_2, X_k] = b_k X_5, \quad 6 \leq k \leq n, \\ [X_i, X_j] = c_{i,j} X_5, \quad 6 \leq i < j \leq n, \\ \text{with } \sum_{i=6}^{j-1} c_{i,j} d_i - \sum_{i=j+1}^{n} c_{j,i} d_i = 0 \text{ for } 6 \leq j \leq n. \end{cases}$$

The first family is a plane of dimension $\frac{(n-4)(n-3)}{2}$. The second family defines an algebraic variety of codimension $n-5$ is a space of dimension $\frac{(n-4)(n-3)}{2}$.

Proposition 8 gives all the nilpotent Lie algebras of breadth 3 and characteristic sequence $(4, 1, \cdots, 1)$. The nilpotent Lie algebras of breadth 3 and characteristic sequences $(3, 2, 1, \cdots, 1)$ and $(2, 2, 2, 1, \cdots, 1)$ are determinated in [8].

Remark. We have just seen that the set $\mathcal{F}^{n,4}_{(1,0,0,n-5)}$ of n-dimensional Lie algebras, $n \geq 5$, whose characteristic sequence is $(4, 1, \cdots, 1)$ is an algebraic variety which is the union of two irreducible components, the first one is the orbit by the linear group of a $\frac{(n-4)(n-3)}{2}$ plane then it is reduced (the affine shema which corresponds to this component is reduced). Then, if there exists a rigid Lie algebra $\mathfrak{g}_1$ in $\mathcal{F}^{n,4}_{(1,0,0,n-5)}$, that is any deformation of $\mathfrak{g}_1$ in $\mathcal{F}^{n,4}_{(1,0,0,n-5)}$ is isomorphic to $\mathfrak{g}_1$, then $\dim H^2_{CR}(\mathfrak{g}_1, \mathfrak{g}_1) = 0$, where $H^2_{CR}(\mathfrak{g}_1, \mathfrak{g}_1) = \frac{Z^2_{CR}(\mathfrak{g}_1, \mathfrak{g}_1)}{B^2_{CR}(\mathfrak{g}_1, \mathfrak{g}_1)}$ and the two cocycles of $Z^2_{CR}(\mathfrak{g}_1, \mathfrak{g}_1)$ are bilinear maps satisfying (6) and $B^2_{CR}(\mathfrak{g}_1, \mathfrak{g}_1)$ the subspace

$\{\delta f, f \in gl(n,\mathbb{K})\} \cap Z^2_{CR}(\mathfrak{g}_1,\mathfrak{g}_1)$. Now, if we consider the Lie algebra

$$\begin{cases} [X_1, X_i] = X_{i+1}, \quad i = 2,3,4, \\ [X_2, X_3] = X_5, \\ [X_2, X_{n-1}] = X_5, \\ [X_2, X_n] = X_4, \\ [X_3, X_n] = X_5, \\ [X_6, X_7] = \cdots = [X_r, X_n] = X_5, \ r = 2p \text{ if } n = 2p+1 \text{ or } r = 2p-1 \text{ if } n = 2p. \end{cases}$$

we have $\dim Z^2_{CR}(\mathfrak{g}_1,\mathfrak{g}_1) = \frac{(n-4)(n-3)}{2} = \dim B^2_{CR}(\mathfrak{g}_1,\mathfrak{g}_1)$. We deduce that $\mathfrak{g}_1$ is rigid in $\mathcal{F}^{n,4}_{(1,0,0,n-5)}$ and the first component is the Zariski closure of the orbit, for the natural action of the linear group, of $\mathfrak{g}_1$. A similar computation for the second component shows that the Lie algebra $\mathfrak{g}_2$ given by

$$\begin{cases} [X_1, X_i] = X_{i+1}, \quad i = 2,3,4, \\ [X_2, X_3] = X_n, \\ [X_2, X_n] = X_5, \\ [X_{2k}, X_{2k+1}] = X_5 \ \ 3 \leq k \leq \frac{n-1}{2}, \end{cases}$$

belongs to the second component and it is rigid in this component. So any n-dimensional Lie algebras of characteristic sequence $(4,1,\cdots,1)$ is a contraction (or degeneration) of one of these two Lie algebras.

3.2. **Lie algebras of characteristic sequence** $(2,2,2,1,\cdots,1)$**.** Let $n \geq 8$ and let $\mathfrak{g}_{3,n-6}$ be the n-dimensional Lie algebra given by the brackets

$$[X_1, X_{2i}] = X_{2i+1}, \ 1 \leq i \leq 3,$$

and other non defined brackets are equal to zero. Any 2-step nilpotent n-dimensional Lie algebra with characteristic sequence $(2,2,2,1,\cdots,1)$ is isomorphic to a linear deformation of $\mathfrak{g}_{3,n-6}$. Its Lie bracket is isomorphic to $\mu = \mu_0 + t\varphi$, where μ_0 is the Lie bracket of $\mathfrak{g}_{3,n-6}$, and φ is a skew-bilinear form such that

$$\begin{cases} \varphi \in Z^2_{CH}(\mathfrak{g}_{3,n-6}, \mathfrak{g}_{3,n-6}), \\ \varphi \circ_1 \varphi = 0. \end{cases}$$

From the construction of the linear deformation, we can assume that $\varphi(X_1, X) = 0$ for any X. Now the first relation is equivalent to

$$\begin{cases} \varphi(X_{2i}, X_{2j}) = \displaystyle\sum_{k=1}^{3} a^{2k+1}_{2i,2j} X_{2k+1} + \sum_{k=8}^{n} a^{k}_{2i,2j} X_k, \ 1 \leq i < j \leq 3, \\ \varphi(X_s, X_l) = \displaystyle\sum_{k=1}^{3} b^{2k+1}_{s,l} X_{2k+1}, \ s, l \geq 8. \end{cases}$$

In this case, the identity $\varphi \circ_1 \varphi = 0$ is satisfied and with such cocycles we describe, up to isomorphism, all the deformations of $\mathfrak{g}_{3,n-6}$ which are 2-step nilpotent. But, we have to restrict these deformations only to Lie algebras with characteristic sequence equal to $(2,2,2,1,\cdots,1)$. This implies that the breadth is equal to 3. This reduces the expression of the cocycles to

(1) $n \geq 9$

$$\begin{cases} \varphi_1(X_2, X_4) = \sum_{k=1}^{3} a_{2,4}^{2k+1} X_{2k+1} + X_8, \\ \varphi_1(X_2, X_6) = \sum_{k=1}^{3} a_{2,6}^{2k+1} X_{2k+1} + X_9, \\ \varphi_1(X_4, X_6) = \sum_{k=1}^{3} a_{4,6}^{2k+1} X_{2k+1}, \\ \varphi_1(X_s, X_l) = 0 \ s, l \geq 8. \end{cases}$$

(2) $n \geq 8$

$$\begin{cases} \varphi_2(X_2, X_4) = \sum_{k=1}^{3} a_{2,4}^{2k+1} X_{2k+1} + X_8, \\ \varphi_2(X_{2i}, X_{2j}) = \sum_{k=1}^{3} a_{2i,2j}^{2k+1} X_{2k+1}, \ 2 \leq i < j \leq 3, \\ \varphi_2(X_s, X_l) = a_{s,l} X_5, \ s, l \geq 8. \end{cases}$$

(3) $n \geq 8$

$$\begin{cases} \varphi_3(X_{2i}, X_{2j}) = \sum_{k=1}^{3} a_{2i,2j}^{2k+1} X_{2k+1}, \ 1 \leq i < j \leq 3, \\ \varphi_3(X_s, X_l) = \sum_{k=1}^{3} a_{s,l} X_{2k+1}, \ s, l \geq 8. \end{cases}$$

Proposition 9. *Any n-dimensional nilpotent Lie algebra whose breadth is 3 and characteristic sequence $(2, 2, 2, 1, \cdots, 1)$ is isomorphic to a Lie algebra whose Lie bracket is $\mu_0 + \varphi_i$ where φ_i, $i = 1, 2, 3$ are defined above.*

For the 7-dimensional case, since the classification of 7-dimensional nilpotent Lie algebras is known, from this list we find the following 7-dimensional nilpotent Lie algebras of characteristic sequence $(2, 2, 2, 1)$ (the notations are those of [**?**])

$\mathfrak{n}_7^{120}$: $[X_1, X_i] = X_{i+1}$, $i = 2, 4, 6$, $[X_2, X_4] = X_7$.

$\mathfrak{n}_7^{121}$: $[X_1, X_i] = X_{i+1}$. $i = 2, 4, 6$

$\mathfrak{n}_7^{122}$: $[X_1, X_i] = X_{i+1}$, $i = 2, 4, 6$, $[X_4, X_6] = X_7$.

$\mathfrak{n}_7^{123}$: $[X_1, X_i] = X_{i+1}$, $i = 2, 4, 6$, $[X_2, X_4] = X_5$, $[X_4, X_6] = X_3$.

3.3. **Lie algebras of characteristic sequence** $(3, 2, 1, \cdots, 1)$**.** Consider now the case of characteristic sequence $(3, 2, 1, \cdots, 1)$.Thus we assume that the dimension of $\mathfrak{g}$ is greater than 6. In dimension 6 any Lie algebra with characteristic sequence $(3, 2, 1)$ is isomorphic to

$\mathfrak{n}_6^{11}$: $[X_1, X_i] = X_{i+1}$, $i = 2, 3, 5$, $[X_5, X_6] = X_4$

$\mathfrak{n}_6^{12}$: $[X_1, X_i] = X_{i+1}$, $i = 2, 3, 5$, $[X_2, X_5] = X_4$

$\mathfrak{n}_6^{13}$: $[X_1, X_i] = X_{i+1}$, $i = 2, 3, 5$, $[X_2, X_3] = X_6$ $[X_2, X_5] = X_6$

$\mathfrak{n}_6^{14}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ [X_2, X_3] = X_4 - X_6\ [X_2, X_5] = X_6$

$\mathfrak{n}_6^{15}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ [X_2, X_5] = X_6\ [X_5, X_6] = X_4$

$\mathfrak{n}_6^{16}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5\ [X_2, X_3] = X_4$

$\mathfrak{n}_6^{17}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5$

In dimension 7, any indecomposable Lie algebra with characteristic sequence $(3, 2, 1, 1)$ is isomorphic to

$\mathfrak{n}_7^{93}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_5] = X_7.$

$\mathfrak{n}_7^{94}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_5] = X_4,\ [X_2, X_3] = X_7$

$\mathfrak{n}_7^{95}$: $\begin{array}{l}[X_1, X_i] = X_{i+1},\ i = 2, 3, 5\\ [X_2, X_3] = X_7\end{array}$

$\mathfrak{n}_7^{96}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_3, X_5] = -X_4,\ [X_2, X_6] = X_4.$

$\mathfrak{n}_7^{97}$: $\left\{\begin{array}{l}[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_6] = X_4,\ [X_3, X_5] = -X_4,\\ [X_2, X_5] = X_7.\end{array}\right.$

$\mathfrak{n}_7^{98}$: $\left\{\begin{array}{l}[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_6] = X_4,\ [X_3, X_5] = -X_4,\\ [X_2, X_5] = X_7,\ [X_5, X_6] = X_4.\end{array}\right.$

$\mathfrak{n}_7^{99}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_7] = X_6,\ [X_2, X_3] = X_4.$

$\mathfrak{n}_7^{100}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_7] = X_4,\ [X_5, X_7] = X_6.$

$\mathfrak{n}_7^{101}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_7] = X_6,\ [X_5, X_7] = X_4.$

$\mathfrak{n}_7^{102}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_5, X_7] = X_4.$

$\mathfrak{n}_7^{103}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_7] = X_4.$

$\mathfrak{n}_7^{104}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_5, X_7] = X_4,\ [X_2, X_3] = X_4.$

$\mathfrak{n}_7^{105}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_7] = X_4,\ [X_2, X_3] = X_4.$

$\mathfrak{n}_7^{106}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_7] = X_4,\ [X_5, X_6] = X_4.$

$\mathfrak{n}_7^{107}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_5, X_7] = X_3,\ [X_6, X_7] = X_4.$

$\mathfrak{n}_7^{108}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_7] = X_6,\ [X_2, X_3] = X_6.$

$\mathfrak{n}_7^{109}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_5, X_7] = X_6,\ [X_2, X_3] = X_6.$

$\mathfrak{n}_7^{110}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_3] = X_6,\ [X_5, X_7] = X_4.$

$\mathfrak{n}_7^{111}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_7] = X_6,\ [X_2, X_5] = X_4.$

$\mathfrak{n}_7^{112}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_3] = X_6,\ [X_2, X_7] = X_4.$

$\mathfrak{n}_7^{113}$: $[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_5, X_7] = X_6,\ [X_5, X_6] = X_4.$

$\mathfrak{n}_7^{114}$: $\left\{\begin{array}{l}[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_7] = X_4,\ [X_5, X_6] = X_4,\\ [X_5, X_7] = X_6.\end{array}\right.$

$\mathfrak{n}_7^{115}$: $\left\{\begin{array}{l}[X_1, X_i] = X_{i+1},\ i = 2, 3, 5,\ \ [X_2, X_5] = X_4,\ [X_5, X_7] = X_3,\\ [X_6, X_7] = X_4.\end{array}\right.$

$$\mathfrak{n}_7^{116}: \begin{cases} [X_1,X_i]=X_{i+1},\ i=2,3,5, & [X_3,X_5]=-X_4,\ [X_2,X_6]=X_4,\\ [X_5,X_7]=-X_4. \end{cases}$$

$$\mathfrak{n}_7^{117}(\alpha): \begin{cases} [X_1,X_i]=X_{i+1},\ i=2,3,5, & [X_2,X_5]=X_7,\ [X_2,X_7]=X_4,\\ [X_5,X_6]=X_4,\ [X_5,X_7]=\alpha X_4. \end{cases}$$

$$\mathfrak{n}_7^{118}: \begin{cases} [X_1,X_i]=X_{i+1},\ i=2,3,5, & [X_2,X_5]=X_7,\ [X_2,X_6]=X_4,\\ [X_3,X_5]=-X_4,\ [X_5,X_7]=-\frac{1}{4}X_4. \end{cases}$$

Let us consider now the general case. We know that any n-dimensional nilpotent Lie algebra with characteristic sequence $(3,2,1,\cdots,1)$ is isomorphic to a linear deformation of $\mathfrak{g}_{1,1,n-5}$ whose Lie bracket μ_0 is given by

$$\mu_0(X_1,X_i)=X_{i+1},\quad i=2,3,5,$$

other non defined product are equal to zero. Since the classification up an isomorphism seems also in this case very utopic (see the previous example of dimension 7), we shall determine a reduced family containing only the algebras with this characteristic sequence. For this it is sufficient to compute the 2-cocycle of the Chevalley cohomology of $\mathfrak{g}_{1,1,n-5}$ satisfying

$$\begin{cases} \delta^2_{C,\mu_0}(\varphi)=0,\\ \varphi\bullet\varphi=0,\\ \delta^2_{R,\mu_0}(\varphi)=0,\\ \mu_0\circ_1\varphi^2+\varphi\circ_1\mu_0\circ_1\varphi+\varphi^2\circ_1\mu_0=0,\\ \varphi^3=0, \end{cases} \tag{8}$$

with $\delta^2_{R,\mu_0}(\varphi)=\mu_0^2\circ_1\varphi+\mu_0\circ_1\varphi\circ_1\mu_0+\varphi\circ_1\mu_0^2=0$. But any linear deformation of μ_0 associated with a such cocycle belong to $\mathcal{F}^{n,3}$ that is the family of 3-step nilpotent Lie algebras and not necessarily in the sub-family $\mathcal{F}^{n,3}_{1,1,n-5}$ of 3-step nilpotent Lie algebras with characteristic sequence $(3,2,1,\cdots,1)$.

Lemma 10. *We have*

$$\mathcal{F}^{n,3}_{1,1,n-5}=\{\mu\in\mathcal{F}^{n,3},\ \text{with breadth}(\mu)=3\}.$$

From the construction of the linear deformation between any Lie algebra of $\mathcal{F}^{n,3}_{1,1,n-5}$ and μ_0, we can assume that $\varphi(X_1,X)=0$ for any X. Now, since $\mu=\mu_0+\varphi\in\mathcal{F}^{n,3}_{1,1,n-5}$, we have necessarily

$$\varphi(X_i,X_j)\in\mathbb{K}\{X_3,X_4,X_6\}.$$

The previous identities and lemma imply that φ satisfies:

$$\begin{cases} \varphi(X_2,X_3)=c_1X_4+e_1X_6+f_{23}X_7,\\ \varphi(X_2,X_5)=c_2X_4+e_2X_6+f_{25}X_7,\\ \varphi(X_2,X_6)=c_3X_4+e_3X_6+f_{26}X_7,\\ \varphi(X_3,X_5)=(-c_3+b_2)X_4+(d_2-e_3X_6)+f_{26}X_7,\\ \varphi(X_5,X_6)=c_4X_4+f_{56}X_7,\\ \varphi(X_2,X_i)=c_{2i}X_4+e_{2i}X_6,\ i\geq 7,\\ \varphi(X_5,X_i)=b_{5i}X_3+c_{5i}X_4+e_{5i}X_6,\ i\geq 7,\\ \varphi(X_6,X_i)=b_{5i}X_4 \end{cases} \tag{9}$$

References

[1] Ancochea-Bermúdez J.M., Gómez-Martin J.R., Valeiras G., Goze M. Sur les composantes irréductibles de la variété des lois d'algèbres de Lie nilpotentes. J. Pure Appl. Algebra 106 (1996), no. 1, 11–22.
[2] Ancochea-Bermúdez J. M., Goze M. Sur la classification des algèbres de Lie nilpotentes de dimension 7. *C. R. Acad. Sci. Paris* 302 (1986), 611–613.
[3] Cabezas J. M., Gómez J. R. (n-4)-filiform Lie algebras. Comm. Algebra 27 (1999), no. 10, 4803-4819.
[4] Cabezas J. M., Gómez J. R., Jimenez-Merchán A. Family of p-filiform Lie algebras. Algebra and operator theory (Tashkent, 1997), 93-102, Kluwer Acad. Publ., Dordrecht, 1998.
[5] Goze M., Khakimdjanov Y. *Nilpotent Lie algebras.* Mathematics and its Applications, 361. Kluwer Academic Publishers Group, Dordrecht, 1996. xvi+336 pp. ISBN: 0-7923-3932-0
[6] Khuhirun B., Misra K.C., Stitzinger E. On nilpotent Lie algebras of small breadth. *Journal of Algebra* 444 (2015) 328-338.
[7] Remm E., Goze M. Symplectic structures on 2-step nilpotent Lie algebras *Arxiv (2015)*
[8] Remm E., Goze M. k-step nilpotent Lie algebras *Georgian Math. J.* 22 (2015), no. 2, 219-234.
[9] Remm E. Filiform Lie algebras. *preprint Mulhouse* 2016.

Université de Haute Alsace, LMIA, 6 rue des Frères Lumière, 68093 Mulhouse
E-mail address: elisabeth.remm@uha.fr